

# OB TRAIN™ for AP/L&D

Transport	CAR (Discharge)	BLS	ALS	SPC	SHELTER IN PLACE
Labor status	None	Early	Cervical dilation ≥4 cm	At risk for en route delivery	If delivery is imminent, or patient is unsafe for transport
Mobility	Ambulatory*	Ambulatory or Non-ambulatory	Non-ambulatory	Non-ambulatory	
Epidural status	None	Placement ≥1 h**	Placement <1 h**	N/A	
Maternal Support	None	Low	Moderate	High	
Fetal monitoring in transit	No	No	Yes	Yes	

#### MOBILITY:

- \*Able to rise from a standing squat

#### EPIDURAL STATUS :

- \*\*Epidural catheter capped off

#### FETAL MONITORING:

- Intermittent/continuous when available
- If need intrauterine resuscitation measures en route, divert to closer facility options

#### MATERNAL SUPPORT:

- **Low: Non-acute conditions requiring continued hospitalization** (e.g. term healthy mother in early labor/spontaneous rupture of membranes not in labor)
- **Moderate: Ongoing acute conditions requiring continued hospitalization and intermittent IV fluid/medications** (e.g. pyelonephritis, preterm premature rupture of membranes, stable chronic hypertension)
- **High: Medically high risk requiring transport by RN/MD and/or continuous IV medication infusion** (e.g. severe preeclampsia/hypertension on magnesium, insulin dependent diabetes in labor)