



## OB TRAIN® for Postpartum

Transport	Car (Discharge)	BLS/Medivan	ALS/CCT	SPC	SHELTER IN PLACE	
Delivery	VD >6 h or CD >48 h	VD <6 h or CD <48 h	Complicated VD or CD	Medically complicated	Patient is unsafe	
Mobility	Ambulatory	Ambulatory with assistance or wheelchair	Stretcher	Specialized equipment		
Post Op	Non-CD surgery >2 h**	>2 h from CD	<2 h from CD	Medically complicated	for transport	
Maternal Risk	None	Low	Moderate	High		

Neonatal	Well baby/	Intermediate	NICU	Domico
status (circle)	rooming in nursery	intermediate	NICO	Demise

VD = Vaginal delivery; CD = Cesarean delivery

- Mobility: \*Able to rise from a standing squat
- \*\* If adult supervision is available for 24 hours

## **Maternal support:**

- Low: Non-acute conditions but requiring continued hospitalization (e.g. observation after complicated vaginal delivery)
- Moderate: Ongoing acute conditions requiring continued hospitalization and intermittent IV fluid/medications (e.g. chorioamnionitis requiring IV antibiotics, hypertension off magnesium, blood transfusion)
- High: Medically high- risk requiring transport by RN/MD and/or continuous IV medication infusion (e.g. preeclampsia with severe features on magnesium, insulin dependent diabetes on IV insulin)